

Signals for Majorana neutrinos in a $\gamma\gamma$ collider

J.Peressutti and O.A.Sampayo

*Departamento de Física, Universidad Nacional de Mar del Plata
Funes 3350, (7600) Mar del Plata, Argentina*

Abstract

We study the possibilities to detect Majorana neutrinos in $\gamma\gamma$ colliders for different center of mass energies. We study the $W^\pm W^\pm l_j^\mp l_k^\mp$ ($l_j \equiv e, \mu, \tau$) final states which are, due to leptonic number violation, a clear signature for intermediate Majorana neutrino contribution. Such a signal (final dileptons of the same sign) is not possible if the heavy neutrinos are Dirac particles. We present our results for the total cross-section as a function of the neutrino mass and the center of mass energies.

I. INTRODUCTION

Massive neutrinos can come in two different types: as Dirac or Majorana particles. Dirac fermions have distinct particles and antiparticle degrees of freedom while Majorana fermions make no such distinction and have half as many degrees of freedom [1]. In this conditions fermions with conserved charges (color, electric charged, lepton number,...) must be of Dirac type, while fermions without conserved charges may be of either type. New neutrinos could have large masses and be of either type. If there are heavy neutrinos (N), then the present and future experiments offer the possibility of establishing their nature. The production of Majorana neutrinos via e^+e^- , $e^-\gamma$, $\gamma\gamma$ and hadronic collision have been extensively investigated in the past [2,3,4,5,6,7,8,9]. In this work we study the possibility of the $\gamma\gamma$ linear collider to produce clear signatures for Majorana neutrinos. The photon linear collider [10] may be the best alternative to the electron positron colliders. In this paper we discuss the possible detection of Majorana neutrinos in the reactions $\gamma\gamma \rightarrow W^\pm W^\pm l_j^\mp l_k^\mp$ ($l_j \equiv e, \mu, \tau$) where the real gamma beam are obtained by Compton backscattering of laser photons off the linear electron beam. In this process most of the photons are produced at the high energy regions. For the cross section calculation we have used the helicity formalism [11,12], considering the complete set of Feynman diagrams show in Fig.1. The phase space integration was done taking account that the intermediate neutrinos can be either on shell or off shell. In this point the present work differ of previous calculation [9] where the restriction $M_N > \sqrt{s}$ is considered and then only neutrinos off-shell are take into account. Of course, for $M_N < \sqrt{s}$ the new heavy neutrino is more likely to be first discovered via single production in $e^+e^- \rightarrow \nu N$. However the $\gamma\gamma$ collider have some facilities that make it suitable to search Majorana neutrinos and study their interactions. In particular a $\gamma\gamma$ collider offer the possibility to control the helicity of the colliding photons controlling the helicity of the Laser photons. Using these polarized high energy photon beams we have the possibility to study in detail the interactions of Majorana neutrinos and reject possible backgrounds. Due the large CM energies of these colliders we have considered that the mass of the final lepton vanishes.

For the couplings of the Majorana neutrinos we follow Ref [6] starting with rather general lagrangian densities for the interaction of N with W and light leptons l_j (e, μ, τ):

$$\mathcal{L}_{NWl} = - \sum_{j=1}^3 \frac{g B_L^{(j)}}{\sqrt{2}} \bar{l}_j \gamma^\mu P_L N W_\mu^- + h.c. \quad (1)$$

The heavy Majorana neutrino couples to the three flavors lepton with couplings proportional to $B_L^{(j)}$, where j labels the family. This $(B_L^{(j)})^2$ constant are, in notation of [13], the sin square of the mixing angle between light and heavy neutrinos $((S_L^{\nu_e})^2, (S_L^{\nu_\mu})^2, (S_L^{\nu_\tau})^2)$. The constant g is the standard $SU(2)_L$ gauge coupling.

This $B_L^{(j)}$ parameters will affect the final results via the combinations

$$H_i = |B_L^{(i)}|^2 \quad \text{and} \quad H = \sum_{j=1}^3 |B_L^{(j)}|^2 \quad (2)$$

The final leptons can be either of e^- , μ^- or τ^- because this is allowed by the interaction lagrangian (eq.1). All these possible final states are a clear signal for intermediary Majorana neutrino and then we sum the cross section over the flavors of the final lepton. This sum depend of H , H_1 , H_2 and H_3 in the numerator of the total cross section. In the other hand this cross-section also depends on H through the total width $\Gamma_{N \rightarrow all}$ in the Majorana neutrino propagator. This last dependence is relevant when the Majorana neutrino is on-shell. We use the Breit-Wigner propagator for the Majorana neutrino N for different values of the mass M_N . The total width $\Gamma_{N \rightarrow all}$ of N was determined at tree level considering the dominant decay modes $N \rightarrow W^\pm l_j^\mp$:

$$\Gamma_{N \rightarrow all} = \frac{g^2 H}{(32\pi M_N^3 M_W^2)} (M_N^2 - M_W^2) (M_N^4 + M_N^2 M_W^2 - 2M_W^4) \quad (3)$$

The constants H_i ($i = 1, 2, 3$) are severely restricted by available experimental data (CERN e^+e^- collider (LEP) and low energies data) [13,14]. This bounds are

$$\begin{aligned} H_1 &< 0.0066 \\ H_2 &< 0.006 \\ H_3 &< 0.018 \end{aligned} \quad (4)$$

In this conditions the upper bound for H is $H < 0.031$. However the bound on H_1 is even more restricted once the constraint from $\beta\beta_{0\nu}$ are taken into account [15]. It is $H_1 < 2 \times 10^{-5}$ for $M_N < 1$ TeV and then the electronic channel do not contribute appreciably in the kinematic region examined in the present work. The above mentioned bound do not constraint the muonic and tauonic channel and do not put new limits on the constants H_2 and H_3 . Then, when the bounds from $\beta\beta_{0\nu}$ are considered the new upper bound on H is

$$H < 0.024. \quad (5)$$

It is the value we have considered in the present article. In this work we have considered the complete set of Feynman diagrams (Fig.1) that contribute at tree level to $\gamma\gamma \rightarrow W^\pm W^\pm l_j^\mp l_k^\mp$ ($\rightarrow jets + l_j l_k$) with the light leptons $l_1 = e, l_2 = \mu, l_3 = \tau$.

In $|\bar{M}|^2$ we average over the initial photon polarizations and sum over the final polarization of the both W and l_j^- and over the flavor of the final leptons. Moreover an $\frac{1}{2}$ factor is included to avoid double counting of the two W and another $\frac{1}{2}$ to avoid double counting of the two identical leptons when integrating over the phase space. We take as inputs the values of $\sqrt{s}$ and M_N . The H dependence is complicate due to the Majorana neutrino propagator. We have founded that the most important contribution to $|\bar{M}|^2$ coming from the couplings is proportional to H^2 . However in the $M_W < M_N < \sqrt{s} - M_W$ kinematic region (Reg.I), where the intermediate Majorana neutrino may be on-shell, the total cross-section is approximately proportional to the H value. In the other hand in the $M_N > \sqrt{s} - M_W$ region (Reg.II), where the Majorana neutrino is off-shell, the total cross-section is approximately proportional to H^2 . The behaviour in Reg.I is easy to realize if we

make the so-called peaking approximation, in which the Breit-Wigner shape of the Majorana neutrino propagator is replaced by a delta function. In this region the H^2 dependence in the numerator is canceled by the H factor in the total width. Considering only the relevant factors in the cross section, we have

$$\sigma = \sum_j \sigma_j \sim \dots H^2 \frac{1}{(q^2 - M_N^2)^2 + M_N^2 \Gamma_N^2} \dots \quad (6)$$

where j labels the final lepton flavors. Making now the peaking approximation

$$\dots \frac{1}{(q^2 - M_N^2)^2 + M_N^2 \Gamma_N^2} \dots \rightarrow \frac{\pi}{M_N \Gamma_N} \delta(q^2 - M_N^2) \quad (7)$$

and since that $\Gamma \sim H$ (eq.3) then we can see that $\sigma = \sum_j \sigma_j$ is approximately proportional to the H value in Reg.I. The expression for M is too long. It extend over tens of page when printed out and we do not present explicitly it here. We show the numerical results for the cross-section for the $\gamma\gamma$ -subprocess for different values of $\sqrt{\hat{s}}$ (the center of mass energie of the $\gamma\gamma$ subprocess) and M_N . The numerical Monte Carlo integration was done using the FORTRAN subroutine RAMBO [16]. The Fig.2 show the M_N dependence of the unpolarized cross-section $\hat{\sigma}$ at fixed $\sqrt{\hat{s}}$ for the maximum value of H given by eq.5. To check the correctness of our final 4-body calculation we include in the same figure the corresponding results for the final 3-body process ($\gamma\gamma \rightarrow l^\pm N W^\pm$), for $\sqrt{\hat{s}} = 500$ GeV. Moreover, partial results were also checked using the CalcHep program [17]. In Fig.3 we show the $\sqrt{\hat{s}}$ dependence of $\hat{\sigma}$ for different values of M_N keeping the same values for H .

In the following the embedding of the subprocess $\gamma\gamma \rightarrow W^\pm W^\pm l_j^\mp l_k^\mp$ into appropriate photon fluxes is presented. We have considered real gamma beam obtained by the Compton backscattering of laser photon off the linear electron beam. The mentioned subprocess is related to e^+e^- collision by folding the $\gamma\gamma$ -subprocess cross-section ($\hat{\sigma}(\hat{s})$) with appropriate $\gamma\gamma$ luminosity function:

$$\sigma(s) = \int_{\tau^{min}}^{\tau^{max}} d\tau \frac{dL_{\gamma\gamma}}{d\tau} \hat{\sigma}_{\gamma\gamma \rightarrow W^\pm W^\pm l_j^\mp l_k^\mp}(\tau s) \quad (8)$$

The differential luminosity $dL_{\gamma\gamma}/d\tau$ is defined as usual in terms of photons structure functions of electrons ($f(x)$) beam:

$$\frac{dL_{\gamma\gamma}}{d\tau} = \int_{\frac{\tau}{x^{max}}}^{x^{max}} \frac{dx}{x} f(x) f\left(\frac{\tau}{x}\right) \quad (9)$$

where $\tau = \hat{s}/s$. $\sqrt{\hat{s}}$ is the center of mass energy of the subsystem and τ^{min} is determined by the production threshold:

$$\tau^{min} = (2m_W)^2/s \quad (10)$$

and x is the fraction of the parent electron's energies carried by the photons. The maximum possible value for x is

$$x^{max} = \frac{\xi}{1 + \xi}, \quad \text{where} \quad \xi = \frac{4E_0\omega_0}{m_e}; \quad (11)$$

where E_0 and ω_0 are the incident electron and laser light energies. To avoid unwanted e^+e^- pair production from the collisions between the incident and back-scattered photons, we should not choose too large ω_0 . This fact constraint the maximum value for ξ to be $\xi = 2(1 + \sqrt{2})$. The variable τ is obviously $\tau = x_a x_b$, the product of electron's energies fractions. In this conditions the minimal value for x is τ/x^{max} and the maximum value for τ is $(x^{max})^2$. This justify the integral limits in eq(8) and (9).

In the other hand we can translate eq(8) and (9) in a alternative expression for the cross-section:

$$\sigma(s) = \int_{x_a^{min}}^{x^{max}} dx_a \int_{\frac{\tau^{min}}{x_a}}^{x^{max}} dx_b f(x_a) f(x_b) \hat{\sigma}(x_a x_b s) \quad (12)$$

where $x_a^{min} = \tau^{min}/x^{max}$.

To illustrate the possible impact of this process in the discovery of Majorana neutrinos we show in Fig 4 the number of events as a function of $\sqrt{s}$ for $M_N = 300 \text{ GeV}$. We have considered the maximum possible value of H as inferred of the experimental bounds (eq.(5)). We have used the estimated luminosity [18] for the $\gamma\gamma$ collider of $\mathcal{L} = 100 \text{ fb}^{-1}$. We have ignored the experimental difficulties of detecting the discussed process unambiguously but if we take as reasonable the threshold of 100 events then we could see signatures for Majorana neutrinos with mass of 300 GeV for center of mass energies greater than 700 GeV. We have also include in Fig.5 a contour plot for different number of events as a function of H and $\sqrt{s}$ for $M_N = 300 \text{ GeV}$. The horizontal line represent the maximum possible value for H .

Summarizing, we calculate the cross-section for the process $\gamma\gamma \rightarrow W^\pm W^\pm l_j^\mp l_k^\mp$ where l_1, l_2 and l_3 are light leptons (e, μ, τ) respectively. We show the total unpolarized cross-section for the subprocess $\gamma\gamma$ for different values of $\sqrt{s}$ and M_N . We have included all the contributions considering that the intermediate Majorana neutrinos can be either on-shell or off-shell. Finally we show the number of events as a function of $\sqrt{s}$ considering the real gamma beam is obtained by the Compton backscattering of laser photons off the linear electron beam.

Acknowledgements

We thank CONICET (Argentina) and Universidad Nacional de Mar del Plata (Argentina) for their financial supports.

REFERENCES

- [1] B. Kayser, F. Gibrat-debu, and F. Perrier, *The Physics of Massive Neutrinos* (World Scientific, Singapore, 1989)
- [2] Ernest Ma and James Pantaleone, Phys. Rev. **D40**, 2172 (1989)
- [3] A. Datta, M. Guchait and A. Pilaftsis, Phys. Rev. **D50**, 3195 (1994)
- [4] J. Gluza and M. Zralek, Phys. Rev. **D51**, 4707 (1995)
- [5] Axel Hoefer and L. M. Sehgal, Phys. Rev **D54**, 1944 (1996)
- [6] G. Cvetcic, C. S. Kim and C. W. Kim, Phys. Rev. Lett. **82**, 1999.
- [7] F. M. L. Almeida Jr, Y. A. Coutinho, J.A. Martins Simoes and M. A. B. do Vale, **hep-ph/0002024**
- [8] J. Peressutti, A.O. Sampayo and J.I. Aranda, Phys. Rev. D **64**, 073007 (2001).
- [9] G. Belanger, F. Bojudeama, D. London and H. Nadeau. Phys. Rev. D **53**, 6292 (1996).
- [10] T. Ohgaki, T. Takahashi and I. Watanabe, Phys. Rev. **D56** 1723 (1997).
- [11] P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, Phys. Lett. **B105**, (1981) 215; P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, Nucl. Phys. **B206**, (1982) 53; F. A. Berends, R. Kleiss, P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, Nucl. Phys. **B206**, (1982) 61; D. Dankaert, P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, Phys. Lett. **B114**, (1982) 203; F. A. Berends, P. De Causmaecker, R. Gastmans, R. Kleiss, W. Troost and T. T. Wu, Nucl. Phys. **B239**, (1984) 382; F. A. Berends, P. De Causmaecker, R. Gastmans, R. Kleiss, W. Troost and T. T. Wu, Nucl. Phys. **B239**, (1984) 395; F. A. Berends, P. De Causmaecker, R. Gastmans, R. Kleiss, W. Troost and T. T. Wu, Nucl. Phys. **B264**, (1986) 234; F. A. Berends, P. De Causmaecker, R. Gastmans, R. Kleiss, W. Troost and T. T. Wu, Nucl. Phys. **B264**, (1986) 265.
- [12] R. Kleiss and W. J. Stirling, Nucl. Phys. **B262** (1985) 235-262.
- [13] E. Nardi, E. Roulet, D. Tomassini, Phys. Lett. **B344**, 225 (1995).
- [14] P. Langacker and D. London, Phys. Rev. D **38**, 886 (1988).
- [15] A. Balysh et al, Phys. Lett. **B356**, 450 (1995).
- [16] R. Kleiss and W. J. Stirling, Comput. Phys. Commun. **40**, 359, 1986
- [17] A. Pukhov et al, "CompHEP - a package for evaluation of Feynman diagrams and integration over multi-particle phase space.", Preprint INP MSU 98-41/542, hep-ph/9908288
- [18] Ilya F. Ginzburg hep-ph/9907549

Figure Captions







Figure 1: Feynman graph contributing to the amplitude of the $\gamma\gamma \rightarrow W^+W^+l^-l^-$ process.

Figure 2: Unpolarized cross-section ($\hat{\sigma}$) for the $\gamma\gamma$ -subprocess as a function of the Majorana neutrino masses (M_N) for different center of mass energies ($\sqrt{\hat{s}}=200, 300$ and 500 GeV). To check the correctness of our final 4-body calculation we include, for $\sqrt{\hat{s}}=500$ GeV, a plot of $\sigma(\gamma\gamma \rightarrow l^\mp NW^\pm)Br(N \rightarrow l^\mp W^\pm)$ (Dot-Dash).

Figure 3: Unpolarized cross section ($\hat{\sigma}$) for the $\gamma\gamma$ -subprocess as a function of the center of mass energies ($\sqrt{\hat{s}}$) for different Majorana neutrino masses ($M_N=200, 300$ and 500 GeV).

Figure 4: Number of events as a function of $\sqrt{s}$ for $M_N = 300$ GeV.

Figure 5: Contour Plot for the number of events (1000, 500, 100, 30, 1) as a function of $\sqrt{s}$ and H . The horizontal line represent the maximum possible value for H .






Fig.1




Fig.2




Fig.3




Fig.4




Fig.5